

BARROW'S INEQUALITY AND SIGNED ANGLE BISECTORS

BRANKO MALEŠEVIĆ, MAJA PETROVIĆ

Submitted to arXiv.org

Abstract. In this paper we give one extension of BARROW's type inequality in the plane of the triangle $\triangle ABC$ introduce signed angle bisectors.

1. Introduction

Let triangle $\triangle ABC$ be given in Euclidean plane. Denote by R_A, R_B and R_C the distances from the arbitrary point M in the plane of $\triangle ABC$ to the vertices A, B and C respectively, and denote by $\ell_a = |MA'|$, $\ell_b = |MB'|$ and $\ell_c = |MC'|$ the length of angle bisectors of $\angle BMC$, $\angle CMA$ and $\angle AMB$ from the point M respectively (Fig. 1).

BARROW's inequality [2]:

$$R_A + R_B + R_C \geq 2(\ell_a + \ell_b + \ell_c) \quad (1)$$

is true when M is arbitrary point in the interior of triangle $\triangle ABC$. The equality holds iff triangle ABC is equilateral and point M is its circumcenter. In this paper we consider a BARROW's type inequality when M is arbitrary point in the plane of the triangle $\triangle ABC$ introduce signed angle bisectors. Let us notice that inequalities with angle bisectors recently are considered in papers [1], [6], [7], [15].




Figure 1: Barrow's inequality (point M into $\triangle ABC$)

Mathematics subject classification (2010): 51M16, 51M04, 14H50.

Keywords and phrases: Signed distance; BARROW's inequality.

Research is partially supported by the Ministry of Science and Education of the Republic of Serbia, Grant No. III 44006 and ON 174032.

Inequality of ERDÖS-MORDELL [4]:

$$R_A + R_B + R_C \geq 2(r_a + r_b + r_c) \quad (2)$$

is a consequence of inequality of BARROW, where r_a , r_b and r_c are distances of interior point M of triangle to the sides BC , CA and AB respectively.

Let us notice that topic of the ERDÖS-MORDELL inequality is current, as it has been shown in recent papers. V. PAMBUCCIAN proved that, in the plane of absolute geometry, the ERDÖS-MORDELL inequality is an equivalent to non-positive curvature [12]. In the paper [11] is given an extension of the ERDÖS-MORDELL inequality on the interior of the ERDÖS-MORDELL curve. In relation to the ERDÖS-MORDELL inequality N. DERGIADIS in the paper [3] proved one extension of the ERDÖS-MORDELL type inequality

$$R_A + R_B + R_C \geq \left(\frac{c}{b} + \frac{b}{c}\right)r'_a + \left(\frac{c}{a} + \frac{a}{c}\right)r'_b + \left(\frac{a}{b} + \frac{b}{a}\right)r'_c \quad (3)$$

where r'_a , r'_b and r'_c are signed distances of arbitrary point M in the plane triangle to the sides BC , CA and AB respectively.

2. The Main Results

Proof of BARROW's inequality in the paper of Z. LU [10] is based on the next theorem.

STATEMENT 1. *Let $p, q, r \geq 0$ and $\alpha + \beta + \gamma = \pi$. Then we have the inequality*

$$p + q + r \geq 2\sqrt{qr} \cos \alpha + 2\sqrt{pr} \cos \beta + 2\sqrt{pq} \cos \gamma. \quad (4)$$

Peculiarity of BARROW's and LU's proofs are, that is, primarily algebraic. In LU's proof, BARROW's inequality follows from positivity of quadratic function $f(x) = x^2 - 2(\sqrt{r} \cos \beta + \sqrt{q} \cos \gamma)x + q + r - 2\sqrt{qr} \cos \alpha$ in the point $x = \sqrt{p}$ with an appropriate geometric interpretation for p, q, r and α, β, γ (for details see [10]).

In this paper we also give one algebraic proof with geometric interpretation for points outside of the triangle $\triangle ABC$. The following theorems are true.

STATEMENT 2. *Let $p, q, r \geq 0$ and $\alpha = \beta + \gamma$. Then we have the inequality*

$$p + q + r \geq -2\sqrt{qr} \cos \alpha + 2\sqrt{pr} \cos \beta + 2\sqrt{pq} \cos \gamma. \quad (5)$$

Proof. Let us consider the quadratic function

$$g(x) = x^2 - 2(\sqrt{r} \cos \beta + \sqrt{q} \cos \gamma)x + q + r + 2\sqrt{qr} \cos \alpha. \quad (6)$$

Then a quarter of the discriminant is

$$\frac{1}{4}\delta = (\sqrt{r} \cos \beta + \sqrt{q} \cos \gamma)^2 - (q + r + 2\sqrt{qr} \cos \alpha). \quad (7)$$

Based on $\alpha = \beta + \gamma$ we have $\cos \alpha = \cos(\beta + \gamma) = \cos \beta \cos \gamma - \sin \beta \sin \gamma$ and hence

$$\begin{aligned} \frac{1}{4} \delta &= r \cos^2 \beta + q \cos^2 \gamma + 2\sqrt{rq} \cos \beta \cos \gamma - q - r - 2\sqrt{rq} \cos \alpha \\ &= r \cos^2 \beta + q \cos^2 \gamma + 2\sqrt{rq} \cos \beta \cos \gamma - q - r - 2\sqrt{rq} \cos(\beta + \gamma) \\ &= -r \sin^2 \beta - q \sin^2 \gamma + 2\sqrt{rq} \cos \beta \cos \gamma - 2\sqrt{rq} \cos \beta \cos \gamma + 2\sqrt{rq} \sin \beta \sin \gamma. \end{aligned}$$

Using previous identity we obtained

$$\delta = -4(\sqrt{r} \sin \beta - \sqrt{q} \sin \gamma)^2 < 0,$$

hence $g(x) \geq 0$. Finally, letting $x = \sqrt{p}$ we obtained (5). $\square$

REMARK 1. Let us emphasize that for term $A = p + q + r + 2\sqrt{qr} \cos \alpha - 2\sqrt{pr} \cos \beta - 2\sqrt{pq} \cos \gamma$, when $\gamma = \alpha - \beta$, follows inequality

$$A = (\sqrt{r} - \sqrt{p} \cos \beta + \sqrt{q} \cos \alpha)^2 + (\sqrt{p} \sin \beta - \sqrt{q} \sin \alpha)^2 \geq 0,$$

analogously using the LAGRANGE's complete square identity from [8], [9]. Therefore we have second proof of inequality (5).

STATEMENT 3. Let $p, q, r \geq 0$ and $\alpha = \beta + \gamma$. Then we have the inequality

$$p + q + r \geq 2\sqrt{qr} \cos \alpha - 2\sqrt{pr} \cos \beta - 2\sqrt{pq} \cos \gamma. \quad (8)$$

Proof. Let us consider the term $A = p + q + r - 2\sqrt{qr} \cos \alpha + 2\sqrt{pr} \cos \beta + 2\sqrt{pq} \cos \gamma$, for $\gamma = \alpha - \beta$. Notice that for the term A , by the LAGRANGE's complete square identity, the following two representations are true.

1 $^\circ$ If $\frac{\pi}{2} \leq \alpha < \pi$, then $\cos \alpha \leq 0$, and therefore

$$A = (\sqrt{r} + \sqrt{p} \cos \beta + \sqrt{q} \cos \alpha)^2 + (\sqrt{p} \sin \beta + \sqrt{q} \sin \alpha)^2 - 4\sqrt{qr} \cos \alpha \geq 0. \quad (9)$$

2 $^\circ$ If $0 < \alpha < \frac{\pi}{2}$, then $\cos \alpha > 0$. From $\alpha = \beta + \gamma$ follows $\cos \beta > 0$, and therefore

$$A = (\sqrt{r} - \sqrt{p} \cos \beta - \sqrt{q} \cos \alpha)^2 + (\sqrt{p} \sin \beta + \sqrt{q} \sin \alpha)^2 + 4\sqrt{pr} \cos \beta \geq 0. \quad \square \quad (10)$$

Let us introduce the division of the plane of triangle $\triangle ABC$ to following areas $\lambda_0 = (+, +, +)$, $\lambda_1 = (-, +, +)$, $\lambda_2 = (+, -, +)$, $\lambda_3 = (+, +, -)$, $\lambda_4 = (+, -, -)$, $\lambda_5 = (-, +, -)$, $\lambda_6 = (-, -, +)$, (Fig. 2), via signes of homogenous barycentric coordinates of a point as given in the paper [14] (see also the Section 7.2 in [5]). Then λ_0 is the interior area of the triangle $\triangle ABC$. Let us notice that $(\lambda_0 \cup \lambda_1) \cup (BC)$ is the interior area of the angle $\angle A$, and λ_4 is the interior area of the opposite angle. Analogously $(\lambda_0 \cup \lambda_2) \cup (AC)$ is the interior area of the angle $\angle B$, λ_5 is the interior area of the opposite angle and $(\lambda_0 \cup \lambda_3) \cup (AB)$ is the interior area of the angle $\angle C$, λ_6 is the interior area of the opposite angle.




Figure 2: The division of the plane by the sidelines of the triangle $\triangle ABC$

The following auxiliary statement is true.

LEMMA 0. Let B and C be fixed points in the plane and let M be arbitrary point in the plane. For ℓ length of angle bisector of $\angle BMC$ from point M following formulas are true:

$$\ell = \frac{2R_B R_C}{R_B + R_C} \cos \frac{\alpha_M}{2} = \frac{\sqrt{R_B R_C}}{R_B + R_C} \sqrt{(R_B + R_C)^2 - |BC|^2}, \quad (11)$$

where $R_B = |MB|$, $R_C = |MC|$ and $\alpha_M = \angle BMC$. Especially, for p line throughout points B and C is true:

$$\ell = \begin{cases} 0 & : M \in [BC], \\ \frac{2R_B R_C}{R_B + R_C} & : M \in p \setminus [BC]. \end{cases} \quad (12)$$

In further considerations let $p = R_A$, $q = R_B$, $r = R_C$. Then, Z. LU, in the paper [10], proved the following BARROW's type inequality.

THEOREM 0. [10] In the area λ_0 the following inequality is true:

$$R_A + R_B + R_C \geq \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) \ell_a + \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) \ell_b + \left(\frac{\sqrt{R_A}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_A}} \right) \ell_c. \quad (13)$$

REMARK 2. BARROW's inequality is a consequence of the previous inequality.

From previous Lemma follows next auxiliary statement.

LEMMA 1. (i) If $M = A$, i.e. $R_A = 0$ then:

$$R_B + R_C \geq \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) \ell_a. \quad (14)$$

(ii) If $M = B$, i.e. $R_B = 0$ then:

$$R_A + R_C \geq \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) \ell_b. \quad (15)$$

(iii) If $M = C$, i.e. $R_C = 0$ then:

$$R_A + R_B \geq \left(\frac{\sqrt{R_B}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_B}} \right) \ell_c. \quad (16)$$

Denote with cl closure of a plane set. The following theorem is true.

THEOREM 1. *In the area $\text{cl}(\lambda_1) \setminus \{B, C\}$ the following inequality is true:*

$$R_A + R_B + R_C \geq \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) (-\ell_a) + \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) \ell_b + \left(\frac{\sqrt{R_A}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_A}} \right) \ell_c. \quad (17)$$

Proof. Let $M \in \text{cl}(\lambda_1) \setminus \{B, C\}$, then $\alpha_M = \beta_M + \gamma_M$ i.e. $\frac{\alpha_M}{2} = \frac{\beta_M}{2} + \frac{\gamma_M}{2}$ (Fig. 3).




Figure 3: Extension of the Barrow's inequality
for the point $M \in \text{cl}(\lambda_1) \setminus \{B, C\}$

Based on the Statement 2, the following inequality holds

$$R_A + R_B + R_C \geq -2\sqrt{R_B R_C} \cos \frac{\alpha_M}{2} + 2\sqrt{R_A R_C} \cos \frac{\beta_M}{2} + 2\sqrt{R_A R_B} \cos \frac{\gamma_M}{2} \quad (18)$$

and based on the Lemma 0 from previous inequality we obtained

$$\begin{aligned} R_A + R_B + R_C &\geq -\frac{R_B + R_C}{\sqrt{R_B R_C}} \ell_a + \frac{R_A + R_C}{\sqrt{R_A R_C}} \ell_b + \frac{R_A + R_B}{\sqrt{R_A R_B}} \ell_c \\ &= \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) (-\ell_a) + \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) \ell_b + \left(\frac{\sqrt{R_A}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_A}} \right) \ell_c. \quad \square \end{aligned} \quad (19)$$

Next two theorems are direct consequence of the Statement 2 by following cyclic replacements $\alpha_M \mapsto \beta_M$, $\beta_M \mapsto \gamma_M$, $\gamma_M \mapsto \alpha_M$ and $R_A \mapsto R_B$, $R_B \mapsto R_C$, $R_C \mapsto R_A$ respectively.

THEOREM 2. *In the area $\text{cl}(\lambda_2) \setminus \{A, C\}$ the following inequality is true:*

$$R_A + R_B + R_C \geq \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) \ell_a + \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) (-\ell_b) + \left(\frac{\sqrt{R_A}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_A}} \right) \ell_c. \quad (20)$$

THEOREM 3. In the area $\text{cl}(\lambda_3) \setminus \{A, B\}$ the following inequality is true:

$$R_A + R_B + R_C \geq \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) \ell_a + \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) \ell_b + \left(\frac{\sqrt{R_A}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_A}} \right) (-\ell_c). \quad (21)$$

The following theorem is true.

THEOREM 4. In the area λ_4 the following inequality is true:

$$R_A + R_B + R_C \geq \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) \ell_a + \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) (-\ell_b) + \left(\frac{\sqrt{R_A}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_A}} \right) (-\ell_c). \quad (22)$$

Proof. Let $M \in \lambda_4$, then $\alpha_M = \beta_M + \gamma_M$ i.e. $\frac{\alpha_M}{2} = \frac{\beta_M}{2} + \frac{\gamma_M}{2}$. Based on the Statement 3 the following inequality is true

$$R_A + R_B + R_C \geq 2\sqrt{R_B R_C} \cos \frac{\alpha_M}{2} - 2\sqrt{R_A R_C} \cos \frac{\beta_M}{2} - 2\sqrt{R_A R_B} \cos \frac{\gamma_M}{2}. \quad (23)$$

Substitutions

$$\ell_a = |MA'| = 2 \frac{R_B R_C}{R_B + R_C} \cos \frac{\alpha_M}{2}, \quad (24)$$

$$\ell_b = |MB'| = 2 \frac{R_A R_C}{R_A + R_C} \cos \frac{\beta_M}{2}, \quad (25)$$

$$\ell_c = |MC'| = 2 \frac{R_A R_B}{R_A + R_B} \cos \frac{\gamma_M}{2} \quad (26)$$

in (23) give

$$\begin{aligned} R_A + R_B + R_C &\geq \frac{R_B + R_C}{\sqrt{R_B R_C}} \ell_a - \frac{R_A + R_C}{\sqrt{R_A R_C}} \ell_b - \frac{R_A + R_B}{\sqrt{R_A R_B}} \ell_c \\ &= \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) \ell_a + \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) (-\ell_b) + \left(\frac{\sqrt{R_A}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_A}} \right) (-\ell_c). \quad \square \end{aligned} \quad (27)$$

Next two theorems are direct consequence of the Statement 3 by following cyclic replacements $\alpha_M \mapsto \beta_M$, $\beta_M \mapsto \gamma_M$, $\gamma_M \mapsto \alpha_M$ and $R_A \mapsto R_B$, $R_B \mapsto R_C$, $R_C \mapsto R_A$ respectively.

THEOREM 5. In the area λ_5 the following inequality is true:

$$R_A + R_B + R_C \geq \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) (-\ell_a) + \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) \ell_b + \left(\frac{\sqrt{R_A}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_A}} \right) (-\ell_c). \quad (28)$$

THEOREM 6. In the area λ_6 the following inequality is true:

$$R_A + R_B + R_C \geq \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) (-\ell_a) + \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) (-\ell_b) + \left(\frac{\sqrt{R_A}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_A}} \right) \ell_c. \quad (29)$$

Now, we give definition of the signed angle bisector for the point M in the plane of the triangle $\triangle ABC$. Let be A fixed vertex and let p be line through vertices B and C . Denote $d = |MA_1|$ distance of the point M to the line p and let $\ell = |MA'|$ be length of the bisector of the angle $\angle BMC$. If d' be signed distance of the point M to the line p related to the vertex A [13] (p. 308.), then $d' = +d$ if M and A with same side of line p , otherwise $d' = -d$. Let us define signed angle bisector ℓ' analogously $\ell' = +\ell$ if M and A with same side of line p , otherwise $\ell' = -\ell$ (Fig. 4). In the case $M \in p$ then $d' = 0$ and then ℓ' given by formula (12).




Figure 4: Signed distances and signed angle bisectors

Let us denote $\mu_1 = \text{cl}(\lambda_1) \setminus \{B, C\}$, $\mu_2 = \text{cl}(\lambda_2) \setminus \{A, C\}$, $\mu_3 = \text{cl}(\lambda_3) \setminus \{A, B\}$, $\mu_4 = \lambda_4$, $\mu_5 = \lambda_5$ and $\mu_6 = \lambda_6$. Then $\bigcup_{i=1}^6 \mu_i \cup \{A, B, C\}$ is a complete division of the plane of the triangle $\triangle ABC$. Finally, analogously to DERGIADES extension of the ERDÖS-MORDELL inequality [3], from previous theorems, an extension of BARROW's type inequality (13) is obtained by the following theorem.

STATEMENT 4. For the point $M \in \bigcup_{i=1}^6 \mu_i$ the following inequality is true:

$$R_A + R_B + R_C \geq \left(\frac{\sqrt{R_C}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_C}} \right) \ell'_a + \left(\frac{\sqrt{R_C}}{\sqrt{R_A}} + \frac{\sqrt{R_A}}{\sqrt{R_C}} \right) \ell'_b + \left(\frac{\sqrt{R_A}}{\sqrt{R_B}} + \frac{\sqrt{R_B}}{\sqrt{R_A}} \right) \ell'_c; \quad (30)$$

otherwise for points $M=A$, $M=B$, $M=C$ following inequalities (14), (15), (16) are true respectively.

REFERENCES

- [1] G.W.I.S.AMARASINGHE, On the standard lengths of Angle Bisectors and Angle Bisector Theorem, Global Journal of Advanced Research on Classical and Modern Geometries, **1**, 1 (2012), 15–27.
- [2] D. F. BARROW, L. J. MORDELL, Solution of Problem 3740, Amer. Math. Monthly, **44**, (1937), 252–254.
- [3] N. DERGIADES, Signed distances and the Erdős-Mordell inequality, Forum Geom., **4**, (2004), 67–68.
- [4] P. ERDÖS, Problem 3740, Amer. Math. Monthly, **42**, (1935), 396.

- [5] R. FAROUKI, *Pythagorean-Hodograph Curves: Algebra and Geometry Inseparable*, Series: Geometry and Computing, Vol. 1, Springer 2008.
- [6] W-D. JIANG, *An Inequality Involving the Angle Bisectors and an Interior Point of a Triangle*, Forum Geometricorum, **8** (2008) 73-76.
- [7] W-D. JIANG AND M. BENCZE, *Some geometric inequalities involving angle bisectors and medians of a triangle*, Journal of Mathematical Inequalities, **5**, 3 (2011), 363-369.
- [8] H. LEE, *Topics in Inequalities - Theorems and Techniques*, Korea Institute for Advanced Study, Seoul, (<http://www.normalesup.org/~kortchem/olympiades/Cours/Inegalites/tin2006.pdf>)
- [9] Z. LU, *Erdős-Mordell inequality and beyond*, Educational Talk at the UCI Math Club (28.11.2007), (<http://www.math.uci.edu/~zlu/talks/2007-uci-mathclub/ucimathclub.pdf>)
- [10] Z. LU, *Erdős-Mordell-type inequalities*, Elemente der Mathematik, **63**, (2008), 23–24.
- [11] B. MALEŠEVIĆ, M. PETROVIĆ, M. OBRADOVIĆ AND B. POPKOSTANTINOVIĆ, *On the Extension of the Erdős-Mordell Type Inequalities*, Mathematical Inequalities and Applications forthcoming articles, MIA-3109, (2013), <http://mia.ele-math.com/forthcoming/>
- [12] V. PAMBUCCIAN, *The Erdős-Mordell inequality is equivalent to non-positive curvature*, Journal of Geometry, **88**, (2008), 134 – 139.
- [13] M. E. SAUL, *Hadamard's Plane Geometry: A Reader's Companion*, American Mathematical Soc., 2010.
- [14] P. YIU, *Conic Construction of a Triangle from the Feet of Its Angle Bisectors*, Journal of Geometry and Graphics, **12**, 2 (2008), 172–182. (<http://math.fau.edu/yiu/YiuFromBisectors.pdf>)
- [15] SH-H. WU AND ZH-H. ZHANG, *A class of inequalities related to the angle bisectors and the sides of a triangle*, J. Inequal. Pure Appl. Math., **7**, 3 (2006), Article 108.

Branko Malešević, Faculty of Electrical Engineering, University of Belgrade,
Bulevar Kralja Aleksandra 73, 11000 Belgrade, Serbia
e-mail: malesevic@etf.rs

Maja Petrović, Faculty of Transport and Traffic Engineering, University of Belgrade,
Vojvode Stepe 305, 11000 Belgrade, Serbia
e-mail: majapet@sf.bg.ac.rs

Corresponding Author: Branko Malešević